


# BENTE 2-row



## Advantages:

- Very good yield
- Excellent yield after grading >2,5 mm
- High diseases resistance; Untreated yield : rel. 109% to french control varieties

## Cultivation:


BENTE is a high yielding barley and a low input type with an ideal and regular protein content for the maltsters.

## Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

### Plant Development



### Vitality



### Plant Health



### Quality



# BENTE 2-row

## Plant development and vitality:

Ear emergence	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Maturity	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

## Yield structure:

Ears/m <sup>2</sup>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Grains/ear	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
TGW	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

## Sowing:

Time of sowing	early until very late
Seed rate	normal conditions, normal time of sowing: 260-280

## Straw:

Lodging resistance	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Necking resistance	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Plant growth regulators	below average, for example 0,2 l/ha Camposan in EC 37-47								

## Disease resistance:

Mildew	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Net blotch	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Rhynchosporium	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Plant protection	Under high disease pressure a splittetd treatment in EC 32 + EC 49 is profitable								

## Quality:

Grading (> 2,2 mm)	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Grading (> 2,5 mm)	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Hectolitre weight	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Raw protein content	low								